

MATHEMATICAL EXPECTATIONS OF FUNCTIONS FROM RANDOM SUMS

Hurshidjon Jumaqulov

Candidate of physical and mathematical sciences, associate professor of the "International tourism and economics" department of the Kokand University

Durdona Bomurotova

Master's student at Karshi State University

Let a sequence of random variables be defined on some probability space (Ω, F, P)

$$\xi_1, \xi_2, \dots, \xi_n, \dots \tag{1}$$

and a non-negative integer random variable ν .

In the theory of summation of random variables, a random sum is a quantity

$$S_\nu = \xi_1 + \xi_2 + \dots + \xi_\nu,$$

i.e. the sum of a random number of random variables (see [4]). In the case where the sequence (1) consists of non-negative independent random variables, the random sum S_ν plays a significant role in the theory of branching random processes ([5], [8]), as well as in reliability theory ([1]). For example, for the study of the asymptotic properties of branching random processes (the probabilities of degeneration and continuation of the process, the limit distributions of the sum of the total number of particles, etc.), B.A. Sevastyanov's problem of describing the class of measurable functions $g(x)$ defined on $[0, \infty)$ such that from the existence of $Eg(\xi_i)$ ($i \in \mathbb{N}$) and $Eg(\nu)$ the existence of mathematical expectation follows $Eg(S_\nu)$.

We first examine the formulated problem for random sums S_ν , where the terms ξ_1, ξ_2, \dots are independent random variables and ν is an independent integer non-negative random variable. According to the formula for the total mathematical expectation, the characteristic function of a random sum S_ν has the following form:

$$\begin{aligned} \Psi(t) = Ee^{itS_\nu} &= \sum_{n=0}^{\infty} E(e^{itS_n}; \nu = n) = \sum_{n=0}^{\infty} P(\nu = n) Ee^{itS_n} = \\ &= \sum_{n=0}^{\infty} P(\nu = n) \prod_{k=1}^n f_k(t), \end{aligned} \tag{2}$$

where $S_0 = 0$, $f_k(t) = Ee^{it\xi_k}$ is the characteristic function of random variables ξ_k . From here on, the entry $E(\xi; A)$ means that the averaging of random variables ξ is performed over a set A , i.e.



$$E(\xi; A) = \int_A \xi(\omega)P(d\omega), \quad A \in F.$$

Taking into account the formulas $ES_\nu^r = i^{-r}\Psi^{(r)}(0)$ from equality (2), we conclude that the existence of a mathematical expectation $\max(E\nu^r, E\xi_i^r) < \infty, i \in \mathbf{N}$, for an integer, $r > 0$ entails $ES_\nu^r < \infty$, and the value ES_ν^r is a function of the values $E\xi_i^{r_1}, E\nu^{r_1}$ for integers r_1 ($0 < r_1 \leq r$). For example, in the case when random variables ξ_i are identically distributed (see [4]),

$$ES_\nu = E\nu E\xi_1, \quad ES_\nu^2 = E\nu[E\xi_1^2 - (E\xi_1)^2] + E\nu^2(E\xi_1)^2$$

etc.

From the above it follows that bounded functions and power functions $g(x) = x^r, r = 1, 2, \dots$ belong to the class of measurable functions possessing the properties formulated in the given problem of B.A. Sevastyanov. Below we will establish fairly broad conditions for functions to belong $g(\cdot)$ to the class of functions considered in this problem.

Following B.A. Sevastyanov [5], we introduce the following classes of functions:

1) We will say that a function $g(x)$ defined on $[0, \infty)$ belongs to the class G_1 if $g(x)$ it is non-negative and there exists such $C > 0$ that for any $x, y \geq 0$

$$g(xy) \leq Cg(x)g(y). \tag{3}$$

2) We will say that a function $g(x)$ defined on $[0, \infty)$ belongs to the class G_2 if it is non-negative, non-decreasing and convex.

The following theorem holds:

Theorem 1 [5]. Let the random variables ξ_i of sequence (1) be independent and identically distributed, and let be ν an independent integer non-negative random variable. Then for any function $g \in G_1 \cap G_2$ the inequality holds.

$$Eg(S_\nu) \leq CEg(\xi_1)Eg(\nu). \tag{4}$$

The monograph [5] presents a proof of Theorem 1 and an application of inequality (4) to the theory of branching random processes. Below we present another proof of this theorem, based on the formula for complete mathematical expectations.

Proof of Theorem 1. Due to convexity $g(x)$ ($g \in G_2$) and from inequality (3) ($g \in G_1$) with probability 1 we have



$$g(S_\nu) = g(\xi_1 + \xi_2 + \dots + \xi_\nu) = g\left(\frac{\xi_1\nu + \xi_2\nu + \dots + \xi_\nu\nu}{\nu}\right) \leq \leq \frac{1}{\nu} \sum_{i=1}^{\nu} g(\xi_i\nu) \leq C \frac{g(\nu)}{\nu} \sum_{i=1}^{\nu} g(\xi_i). \tag{5}$$

Further, according to the formula for the total mathematical expectation and by virtue of inequality (5)

$$E[g(S_\nu)] = \sum_{n=1}^{\infty} E[g(S_\nu); \nu = n] = \sum_{n=1}^{\infty} E[g(S_n); \nu = n] \leq \leq C \sum_{n=1}^{\infty} \left(\frac{g(n)}{n} \sum_{k=1}^n E[g(\xi_k); \nu = n] \right). \tag{6}$$

Considering the identical distribution of random variables ξ_i and their independence from ν , we have

$$E[g(\xi_k); \nu = n] = P(\nu = n) Eg(\xi_1). \tag{7}$$

From relations (6), (7) we finally obtain

$$E[g(S_\nu)] \leq CE[g(\xi_1)] \sum_{n=1}^{\infty} g(n)P(\nu = n) = CE[g(\xi_1)]E[g(\nu)].$$

The proof of Theorem 1 is complete.

Remark 1. As shown by B.A. Sevastyanov [5], the conditions for the existence $Eg(\nu) < \infty$ and membership of a function g in the class of convex functions G_2 cannot be weakened.

Remarks 2. Functions from the class G_1 with a constant $C = 1$ are called semi-multiplicative. From the proof of Theorem 1, we can conclude that if the convexity conditions of functions $g(\cdot)$ are replaced by the concavity condition, then lower bounds for can be obtained $Eg(S_\nu)$. It should also be noted that semi-multiplicative convex functions play an important role in the theory of linear operators (see [2], [3]).



References

1. Гнеденко Б.В., Беляев Ю.К., Соловьев А.Д. Математические методы в теории надежности. Основные характеристики надежности и их статистический анализ. –М. “Наука”, 1965. –524 с.
2. Красносельский М.А., Рудицкий Я.Б. Выпуклые функции и пространства Орлича. – М.: Физматгиз, 1958. – 393 с.
3. Крейн С.Г., Петунин Ю.И., Семенов Е.М. Интерполция линейных операторов. – М.: Наука, 1978. –399 с.
4. Круглов В.М., Королев В.Ю. Предельные теоремы для случайных величин. Изд-во Московского Университета. 1990. –269 с.
5. Севастьянов Б.А. Ветвящиеся процессы. – М.: Наука, 1971. –436 с.
6. Жумакулов Х. К., Каримова Ш. Я. К. НЕКОТОРЫЙ ЗАДАЧИ ВЕТВЯЩИХСЯ ПРОЦЕССОВ //Science and innovation. – 2024. – Т. 3. – №. Special Issue 50. – С. 28-33.
7. Zhumakulov, H. K. "Conditions for the convergence of branching processes with immigration starting from a large number of particles." Galaxy International Interdisciplinary Research Journal 10.12 (2022): 309-313.
8. Rahimov I. Random Sums and Branching stochastic processes. –New York: Springer-Verlag, 1995. –199 p.

